

## 6 • Energy and Chemical Reactions

### P R A C T I C E T E S T

- A system has an increase in internal energy,  $\Delta E$ , of 40 kJ. If 20 kJ of work,  $w$ , is done on the system, what is the heat change,  $q$ ?  
 a) +60 kJ                      d) -20 kJ  
 b) +40 kJ                      e) -60 kJ  
 c) +20 kJ
- A gas at 20 atm pressure with a volume of 2.0 Liters expands against a 5 atm pressure to a volume of 8.0 Liters. How much work is done by the gas?  
 a) 30 L-atm                      c) 8 L-atm  
 b) 18 L-atm                      d) 5 L-atm
- Which equation represents the heat of formation,  $\Delta H_f^\circ$ , for  $\text{MgCl}_2$ ?  
 a)  $\text{Mg}^{2+}(\text{aq}) + 2 \text{Cl}^- \rightarrow \text{MgCl}_2(\text{s})$   
 b)  $\text{Mg}(\text{s}) + 2 \text{Cl}(\text{g}) \rightarrow \text{MgCl}_2(\text{s})$   
 c)  $\text{MgCl}_2(\text{s}) \rightarrow \text{Mg}^{2+}(\text{aq}) + 2 \text{Cl}^-(\text{aq})$   
 d)  $\text{Mg}(\text{s}) + \text{Cl}_2(\text{g}) \rightarrow \text{MgCl}_2(\text{s})$
- Take a toy balloon. Quickly stretch the balloon and press it against your lower lip. What is the  $\Delta H$  for the reaction:  
 unstretched ® stretched  
 a) +                                  c) -  
 b) 0                                  d) impossible to tell
- Which of the following is NOT a state function?  
 a) pressure                      c) temperature  
 b) volume                        d) none of these
- The correct units for specific heat capacity:  
 a)  $\text{J}/^\circ\text{C}$                                   c)  $\text{J/g } ^\circ\text{C}$   
 b)  $\text{J/g}$                                       d)  $^\circ\text{C/g}$
- How much heat is required to convert solid sulfur to gaseous sulfur at 298 K and 1 atm pressure?  

$$\Delta H^\circ (\text{kJ/mol})$$

$\text{S}(\text{s}) + \text{O}_2(\text{g}) \rightarrow \text{SO}_2(\text{g})$	-395
$\text{S}(\text{g}) + \text{O}_2(\text{g}) \rightarrow \text{SO}_2(\text{g})$	-618

 a) -1013 kJ/mol                      c) +223 kJ/mol  
 b) -223 kJ/mol                      d) +618 kJ/mol
- Using the  $\Delta H_f^\circ$  given below, calculate the  $\Delta H_{\text{combustion}}$  for propane,  $\text{C}_3\text{H}_8$ .  

$$\Delta H_f^\circ (\text{kJ/mol})$$

$\text{H}_2\text{O}(\text{l})$	-286
$\text{CO}_2(\text{g})$	-394
$\text{C}_3\text{H}_8(\text{g})$	-104

 a) 576 kJ                                  c) -2222 kJ  
 b) -576 kJ                                d) -2330 kJ
- The heat of vaporization of methane,  $\text{CH}_4$ , at its boiling point is 9.20 kJ/mol. How much heat energy is required to vaporize 100. g of methane at its boiling point?  
 a) 1380 kJ                                  c) 21.6 kJ  
 b) 86.3 kJ                                  d) 57.4 kJ
- How much energy is required to melt 10.0 g benzene,  $\text{C}_6\text{H}_6$ ? The heat of fusion of benzene is 2.37 kJ/mol.  
 a) 3.30 kJ                                  c) 1850 kJ  
 b) 23.7 kJ                                  d) 0.303 kJ

11. If  $\Delta H$  for a reaction is positive, ...
- the reaction rate is generally very fast.
  - the enthalpy change of the reverse reaction is positive.
  - the enthalpy of the products is greater than the enthalpy of the reactants.
  - the energy released during bond formation is greater than the energy absorbed during bonding breaking for the reaction.

12. Given the two equations:



What is the standard enthalpy of formation for sulfur dioxide,  $\text{SO}_2\text{(g)}$ ?

- +99.1 kJ
  - 296.1 kJ
  - 592.2 kJ
  - 839.5 kJ
13. When 0.100 g benzoic acid ( $\text{HC}_6\text{H}_4\text{CO}_2$ ) and excess oxygen is ignited in a bomb calorimeter, the temperature of the water changes from  $25.000^\circ\text{C}$  to  $25.225^\circ\text{C}$ . The heat capacity of the calorimeter is  $603 \text{ J}^\circ\text{C}$ . What is the  $\Delta E$  for this reaction?
- 597 J
  - 1660 J
  - 136 J
  - 149 J
14. Under conditions of constant volume, the heat change that occurs during a chemical reaction is equal to
- $\Delta H$
  - $\Delta E$
  - $\Delta T$
  - $\Delta P$
15. Systems tend toward:
- maximum entropy and minimum enthalpy
  - maximum entropy and maximum enthalpy
  - minimum entropy and minimum enthalpy
  - minimum entropy and maximum enthalpy

### Answers:

- |      |       |       |
|------|-------|-------|
| 1. C | 6. C  | 11. C |
| 2. A | 7. C  | 12. B |
| 3. D | 8. C  | 13. C |
| 4. C | 9. D  | 14. B |
| 5. D | 10. D | 15. A |

### Notes:

- an increase in internal energy means an increase in P.E. of system by 40 kJ  
work done ON system increases P.E., +20 kJ, so  $q = +20 \text{ kJ}$ , too.
- work =  $P\Delta V = 5 \text{ atm} \times (8-2 \text{ L})$ ... the 20 atm is not used for anything.
- balloon gets warm,  $\Delta H < 0$
- each of these only depends on the STATE of the substance, not on its HISTORY.
- reverse second reaction
- recall:  $\text{C}_3\text{H}_8 + 5\text{O}_2 \rightarrow 3\text{CO}_2 + 4\text{H}_2\text{O}$  and use Hess's Law
- Given: 100g  $\text{CH}_4$ , use molar mass &  $H_{\text{vap}}$  as conversion factors.
- Given: 10.0g  $\text{C}_6\text{H}_6$ , use molar mass &  $H_{\text{fus}}$  as conversion factors.
- this is an "uphill" reaction.
- take half of first equation, reverse second equation. if you reverse and double second equation, you get TWICE the answer.
- answer = heat capacity  $\times \Delta T$ ... you don't use 0.100 g anywhere. You would IF the question asked for MOLAR heat of combustion.
- if volume is constant,  $P\Delta V$  work = 0 so  $\Delta E = q + w$  becomes  $\Delta E = q$ .
- from lecture... THIS chapter, however, concentrates on enthalpy,  $\Delta H$ .