

## 6 • Energy and Chemical Reactions

---

### P R A C T I C E T E S T

- How many joules are equivalent to 37.7 cal?  
a) 9.01 J                      c) 1.51 J  
b) 4.184 J                     d) 158 J
- The quantity of heat that is needed to raise the temperature of a sample of a substance 1.00 degree is called its  
a) heat capacity              c) enthalpy  
b) specific heat                d) kinetic energy
- Equal masses of two substances, A & B, each absorb 25 Joules of energy. If the temperature of A increases by 4 degrees and the temperature of B increases by 8 degrees, one can say that  
a) the specific heat of A is double that of B.  
b) the specific heat of B is double that of A.  
c) the specific heat of B is negative.  
d) the specific heat of B is triple that of A.
- If 25 J are required to change the temperature of 5.0 g of substance A by 2.0°C, what is the specific heat of substance A?  
a) 250 J/g°C                    c) 10. J/g°C  
b) 63 J/g°C                     d) 2.5 J/g°C
- How much energy is required to change the temperature of 2.00 g aluminum from 20.0°C to 25.0°C? The specific heat of aluminum is 0.902 J/g°C.  
a) 2.3 J                              c) 0.36 J  
b) 9.0 J                              d) 0.090 J
- Consider the thermal energy transfer during a chemical process. When heat is transferred to the system, the process is said to be \_\_\_\_\_ and the sign of  $\Delta H$  is \_\_\_\_\_.  
a) exothermic, positive  
b) endothermic, negative  
c) exothermic, negative  
d) endothermic, positive
- What is the  $\Delta E$  for a system which has the following two steps:  
Step 1: The system absorbs 60 J of heat while 40 J of work are performed on it.  
Step 2: The system releases 30 J of heat while doing 70 J of work.  
a) 100 J                              c) 30 J  
b) 90 J                                d) zero
- When two solutions react the container “feels hot.” Thus,  
a) the reaction is endothermic.  
b) the reaction is exothermic.  
c) the energy of the universe is increased.  
d) the energy of both the system and the surroundings is decreased.

9. The equation for the standard enthalpy of formation of  $\text{N}_2\text{O}_3$  is
- $\text{N}_2\text{O}(\text{g}) + \text{O}_2(\text{g}) \rightarrow \text{N}_2\text{O}_3(\text{g})$
  - $\text{N}_2\text{O}_5(\text{g}) \rightarrow \text{N}_2\text{O}_3(\text{g}) + \text{O}_2(\text{g})$
  - $\text{NO}(\text{g}) + \text{NO}_2(\text{g}) \rightarrow \text{N}_2\text{O}_3(\text{g})$
  - $\text{N}_2(\text{g}) + \frac{3}{2} \text{O}_2(\text{g}) \rightarrow \text{N}_2\text{O}_3(\text{g})$
10. For the general reaction
- $$2 \text{A} + \text{B}_2 \rightarrow 2 \text{AB}, \quad \Delta H \text{ is } +50.0 \text{ kJ.}$$
- We can conclude that
- the reaction is endothermic.
  - the surroundings absorb energy.
  - the standard enthalpy of formation of AB is  $-50.0 \text{ kJ}$ .
  - the molecule AB contains less energy than A or  $\text{B}_2$ .
11. Calculate the enthalpy of combustion of  $\text{C}_3\text{H}_6$ :
- $$\text{C}_3\text{H}_6(\text{g}) + \frac{9}{2} \text{O}_2(\text{g}) \rightarrow 3\text{CO}_2 + 3\text{H}_2\text{O}$$
- using the following data:
- $$3\text{C}(\text{s}) + 3\text{H}_2(\text{g}) \rightarrow \text{C}_3\text{H}_6(\text{g}) \quad \Delta H^\circ = 53.3 \text{ kJ}$$
- $$\text{C}(\text{s}) + \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) \quad \Delta H^\circ = -394 \text{ kJ}$$
- $$\text{H}_2(\text{g}) + \frac{1}{2} \text{O}_2(\text{g}) \rightarrow \text{H}_2\text{O}(\text{l}) \quad \Delta H^\circ = -286 \text{ kJ}$$
- $-1517 \text{ kJ}$
  - $1304 \text{ kJ}$
  - $-626 \text{ kJ}$
  - $-2093 \text{ kJ}$
12. Which one of the following would have an enthalpy of formation value ( $\Delta H_f^\circ$ ) of zero?
- $\text{H}_2\text{O}(\text{g})$
  - $\text{O}(\text{g})$
  - $\text{H}_2\text{O}(\text{l})$
  - $\text{O}_2(\text{g})$
13. Calculate the heat of vaporization of titanium (IV) chloride:  $\text{TiCl}_4(\text{l}) \rightarrow \text{TiCl}_4(\text{g})$  using the following enthalpies of reaction:
- $$\text{Ti}(\text{s}) + 2\text{Cl}_2(\text{g}) \rightarrow \text{TiCl}_4(\text{l}) \quad \Delta H^\circ = -804.2 \text{ kJ}$$
- $$\text{TiCl}_4(\text{g}) \rightarrow 2\text{Cl}_2(\text{g}) + \text{Ti}(\text{s}) \quad \Delta H^\circ = 763.2 \text{ kJ}$$
- $-1567 \text{ kJ}$
  - $-783.7 \text{ kJ}$
  - $1165 \text{ kJ}$
  - $41 \text{ kJ}$
14. Calculate the enthalpy of reaction for:
- $$\text{D} + \text{F} \rightarrow \text{G} + \text{M}$$
- using the following equations and data:
- $$\text{G} + \text{C} \rightarrow \text{A} + \text{B} \quad \Delta H^\circ = +277 \text{ kJ}$$
- $$\text{C} + \text{F} \rightarrow \text{A} \quad \Delta H^\circ = +303 \text{ kJ}$$
- $$\text{D} \rightarrow \text{B} + \text{M} \quad \Delta H^\circ = -158 \text{ kJ}$$
- $-132 \text{ kJ}$
  - $-422 \text{ kJ}$
  - $+422 \text{ kJ}$
  - $+132 \text{ kJ}$
15. Calculate the standard enthalpy of the reaction for the process
- $$3\text{NO}(\text{g}) \rightarrow \text{N}_2\text{O}(\text{g}) + \text{NO}_2(\text{g})$$
- using the standard enthalpies of formation (in  $\text{kJ/mol}$ ):  $\text{NO} = 90$ ;  $\text{N}_2\text{O} = 82.1$ ;  $\text{NO}_2 = 34.0$
- $-153.9 \text{ kJ}$
  - $206 \text{ kJ}$
  - $-26.1 \text{ kJ}$
  - $386 \text{ kJ}$
16. The standard molar enthalpy of combustion is  $-1277.3 \text{ kJ}$  for the combustion of ethanol.
- $$\text{C}_2\text{H}_5\text{OH}(\text{l}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + 3\text{H}_2\text{O}(\text{g})$$
- Calculate the standard molar enthalpy of formation for ethanol based on the following standard enthalpies of formation:
- $$\Delta H_f^\circ \text{CO}_2 = -393.5 \text{ kJ/mol}$$
- $$\Delta H_f^\circ \text{H}_2\text{O} = -241.8 \text{ kJ/mol}$$
- $-642.7 \text{ kJ/mol}$
  - $-235.1 \text{ kJ/mol}$
  - $235.1 \text{ kJ/mol}$
  - $642.7 \text{ kJ/mol}$

17. Calculate the amount of heat needed to change 25.0 g ice at 0°C to water at 0°C.

The heat of fusion of H<sub>2</sub>O = 333 J/g;

- a) 56.5 kJ                      c) 7.06 kJ  
 b) 8.33 kJ                      d) 463 kJ

**Answers:** (Please use CAPITAL letters) - **V1**

1.   
 2.   
 3.   
 4.   
 5.

11.   
 12.   
 13.   
 14.   
 15.

**Questions 18-20: (1/2 point each)**

The following data was collected in an experiment similar to the Specific Heat experiment performed in class. Fill in the missing values. (Assume the calorimeter has a calorimeter constant of 0 J/°C)

6.   
 7.   
 8.   
 9.   
 10.

16.   
 17.

18.   
 18.   
 19.   
 19.   
 20.   
 20.

Data & Calculations		Glass Beads
	mass of glass beads	4.88 g
	mass of water	10.14 g
	initial temperature of water	19.2 °C
	initial temperature of beads	89.2 °C
	final temperature of mixture	24.8 °C
18.	temp change of water (°C)	
18.	temp change of hot beads (°C)	
19.	change in energy of water, q <sub>water</sub> (J)	
	change in energy of calorimeter (J)	0 J
19.	change in energy of beads, q <sub>beads</sub> (J)	
20.	specific heat of beads (J·g <sup>-1</sup> ·°C <sup>-1</sup> )	
	accepted value of specific heat	.833
20.	% error	