


18 • Acid-Base Reactions

CALCULATIONS


Information from the Curve:

There are several things you can read from the titration curve itself. Consider this titration curve.



- This is a _____ (strong/weak) acid titrated with a strong base. The acid is _____ (monoprotic/diprotic). How would the other strength of acid look?
- Place a dot (●) on the curve at the equivalence point. The pH at the equivalence point is _____. Choose a good indicator for this titration from Figure 17.11 on page 810 of your textbook.
- What volume of base was used to titrate the acid solution? _____ mL
- Place a box (■) on the curve where the pH of the solution = the pK_a of the acid.
 What is the pH at this point? _____
 What is the pK_a of the acid? _____
 What is the K_a of the acid? _____

Calculations knowing the Acid:



- Hydrofluoric acid, HF, has a $K_a = 7.2 \times 10^{-4}$. Calculate the pH of 10.0 mL of a 0.050 M solution of HF. Plot this point on the axes.
- A 0.020 M solution of NaOH is used for the titration. What volume will be needed to reach the equivalence point?
- Write the net reaction for the neutralization of a solution of HF with a solution of NaOH.

- Calculate the moles of F^- at the equivalence point. What is the total volume? _____ L
 The $[F^-]$ at the equivalence point is _____
- Calculate the pH of the solution at the equivalence point. Use this information and the answer to question 6 to plot the equivalence point on your graph. Choose a good indicator for this titration from Figure 17.11 on page 810 of your textbook.

10. What is the pH halfway to the equivalence point? Plot this point on your graph.
11. How many moles of HF are in the original 10.0 mL sample of HF? _____
12. When only 5.0 mL of 0.020 M NaOH has been added, calculate the moles of HF left and F⁻ produced.

	HF	OH ⁻	H ₂ O	F ⁻
<i>i</i>			-----	
<i>c</i>			-----	
<i>e</i>			-----	

13. Use the Henderson-Hasselbach equation or an icebox to calculate the pH when 5.0 mL of base has been added. Plot this point on your graph.
14. When 20.0 mL of 0.020 M NaOH has been added, calculate the moles of HF left and F⁻ produced.

	HF	OH ⁻	H ₂ O	F ⁻
<i>i</i>			-----	
<i>c</i>			-----	
<i>e</i>			-----	


15. Use the Henderson-Hasselbach equation or an icebox to calculate the pH when 20.0 mL of base has been added. Plot this point on your graph.
16. When 30.0 mL of base is added, how many moles of OH⁻ is in excess? _____
 The total volume is _____ L.
 [OH⁻] = _____
 pOH = _____ pH = _____
 Plot this point on your graph.

17. Sketch the titration curve on your graph.

Weak Base-Strong Acid Curve:

A 20.0 mL sample of 0.10 M CH₃NH₂ (methyl amine) is titrated with 0.15 M HCl. The K_b for CH₃NH₂ = 4.2 x 10⁻⁴.

Do the appropriate calculations to sketch a titration curve for this titration.



Formulas from the AP Exam:

EQUILIBRIUM

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

$$K_b = \frac{[\text{OH}^-][\text{HB}^+]}{[\text{B}]}$$

$$K_w = [\text{OH}^-][\text{H}^+] = 1.0 \times 10^{-14} \text{ @ } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log [\text{H}^+], \text{ pOH} = -\log [\text{OH}^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{HB}^+]}{[\text{B}]}$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

$$K_p = K_c(RT)^{\Delta n},$$

where Δn = moles product gas – moles reactant gas