

# Ch 19 – Solubility Equilibria

## Selected NChO Problems

1999-4.	<p><b>(A) AlCl<sub>3</sub></b>            Three clues lead to choice (A):            (1) “colorless aqueous solution” eliminates blue CuSO<sub>4</sub>.            (2) ppt with NaOH eliminates Ba(NO<sub>3</sub>)<sub>2</sub> because Ba(OH)<sub>2</sub> is soluble.            (3) white ppt with AgC<sub>2</sub>H<sub>3</sub>O<sub>2</sub> AgI is a yellow precipitate; AgCl is a white precipitate.</p>
1999-40.	<p><b>(C) 7.3 x 10<sup>-14</sup></b>            PbCO<sub>3</sub> is a 1:1 compound so <math>K_{sp} = s^2</math>  <math>s = 2.7 \times 10^{-7}</math>  <math>K_{sp} = (2.7 \times 10^{-7})^2 = 7.3 \times 10^{-14}</math></p>
1998-4.	<p><b>(D) PbCl<sub>2</sub></b>            Solubility rules help eliminate (A), (B), and (C). (NO<sub>3</sub><sup>-</sup>, Na<sup>+</sup>, and K<sup>+</sup> are always soluble.)</p>
1998-39.	<p><b>(D) 8.8 x 10<sup>-9</sup></b>            PbI<sub>2</sub> is a 1:2 compound so <math>K_{sp} = 4s^3</math>  <math>s = 0.0013 \text{ M}</math> <math>K_{sp} = 4(0.0013)^3 = 8.8 \times 10^{-9}</math></p>
1997-5	<p><b>(C) MgSO<sub>4</sub> and Pb(NO<sub>3</sub>)<sub>2</sub></b>            PbSO<sub>4</sub> precipitate is formed. (solubility rules)</p>
1997-40.	<p><b>(A) 1 only</b>  <math>\text{Ca(OH)}_2 + 2 \text{NH}_4\text{Cl} \rightarrow 2 \text{NH}_3(\text{g}) + 2 \text{H}_2\text{O} + \text{CaCl}_2</math>            “1” is true            “2” CaCl<sub>2</sub> is formed, but it is not a precipitate            “3” is a complex ion.            NH<sub>3</sub> is a good ligand, but you would need concentrated NH<sub>3</sub> to form the complex ion.</p>
1996-40.	<p><b>(B) BaCO<sub>3</sub> only</b>            Look for salt with anion from a weak acid.            BaCO<sub>3</sub>; CO<sub>3</sub><sup>2-</sup> from H<sub>2</sub>CO<sub>3</sub> (weak)            BaSO<sub>4</sub>; SO<sub>4</sub><sup>2-</sup> from H<sub>2</sub>SO<sub>4</sub> (strong)</p>
1995-12.	<p><b>(B) CaCl<sub>2</sub> + K<sub>2</sub>CO<sub>3</sub> ®</b>            CaCO<sub>3</sub> is a precipitate (solubility rules)</p>
1995-35.	<p><b>(C) 4.4 x 10<sup>-4</sup></b>            CuCl is a 1:1 compound, so <math>K_{sp} = s^2</math>  <math>K_{sp} = 1.9 \times 10^{-7} = s^2</math>  <math>s = \sqrt{1.9 \times 10^{-7}} = 4.4 \times 10^{-4}</math></p>
1994-36.	<p><b>(B) 1.5 x 10<sup>-3</sup></b>            PbI<sub>2</sub> is a 1:2 compound so <math>K_{sp} = 4s^3</math>  <math>K_{sp} = 1.4 \times 10^{-8}</math>  <math>s = \sqrt[3]{K_{sp}/4} = 1.5 \times 10^{-3}</math></p>
1993-30.	<p><b>(A) BaCO<sub>3</sub></b>            see explanation for 1996-40. above. HCl, HNO<sub>3</sub>, H<sub>2</sub>SO<sub>4</sub> are all strong acids.</p>
1992-32.	<p><b>(D) CaCl<sub>2</sub></b>            Look for salt with anion NOT from a weak acid. HF, H<sub>2</sub>CO<sub>3</sub>, and H<sub>2</sub>C<sub>2</sub>O<sub>4</sub> are all weak acids.</p>